# IgG 2000 CWP™

# **Immunoglobulin Concentrate from Colostral Whey Peptides**



Available in 30 and 90 servings powder and 120 and 360 capsules

## **Discussion**

Beyond Colostrum: Immunoglobulin Concentrate from Colostral Whey Peptides IgG 2000 CWP is the result of advanced coagulation and filtration technologies that separate bioactive substances and then concentrate them. These precise systems produce a potent, pure, and generally recognized as safe (GRAS) immunoglobulin concentrate from colostral whey peptides. IgG 2000 CWP provides immunoglobulins, including a minimum of 40% IgG; bioactive and growth factors; oligosaccharides; and gangliosides. Each of these components provides the user with different and

#### **Immunoglobulins**

complementary health benefits (see Chart).\*[1]

Oral consumption of immunoglobulins derived from colostrum is a means of supporting passive immunity.<sup>[2-5]</sup> Immunoglobulins flag antigens for the immune system. In doing so, immunoglobulins become key participants in protecting the body and eliminating unwanted molecules. IgG 2000 CWP delivers immunoglobulins and is particularly high (minimum of 40%) in IgG. Among immunoglobulins, IgG is said to be the most versatile, carrying out all of the functions of immunoglobulin molecules.\*

#### **Bioactive and Growth Factors**

Sialic acid is an essential component of mucins, glycoproteins, oligosaccharides, and gangliosides and is therefore important to the function of cell membranes and membrane receptors. It is also important for normal brain development. Sialic acid-containing oligosaccharides in bovine colostrum can prevent certain antigens from binding to host tissues. Lactoferrin is an immune-supporting, iron-binding glycoprotein naturally found in bovine colostrum. It plays an important role in immune regulation and in the body's defense mechanisms. [6] Studies suggest that growth factors from bovine colostrum, including IGF-1 (insulin-like growth factor) and TGF (transforming growth factor), stimulate cell growth in the gut to strengthen the gut lining, help build lean muscle mass, and slow protein catabolism. [1] Furthermore, improvements observed in exercise performance by athletes taking colostrum have been attributed to growth factors.\*

#### **Proline-Rich Peptides**

Proline-rich peptides (PRPs), which were first isolated from ovine colostrum and later from bovine colostrum, are intercellular signaling molecules. They act as regulatory

# **Clinical Applications**

- » Supports Immune Function by Providing Immunoglobulins and Other Immune Factors\*
- » Supports the Body's Normal Gut Repair Pathways\*
- » Helps Maintain a Healthy Microbial and Cytokine Balance in the Gut\*
- » Promotes Overall Health and Well-Being\*

IgG 2000 CWP™ is an immunoglobulin concentrate derived from colostral whey peptides. It delivers natural immunoglobulins (standardized to a minimum of 40% lgG), bioactive proteins, and growth factors. These components support immune function, healthy cytokine activity, gut barrier function, and gastrointestinal health and tissue repair. Advanced coagulation and filtration techniques make lgG 2000 CWP a unique, GRAS formula that is superior in its bioactive composition and its purity.\*

substances that have the unique ability to modulate and stabilize various biologic processes in the body, such as cytokine and immune processes. [1] The in vitro and in vivo effects of PRPs on immunoregulation, including their effects on the maturation and differentiation of thymocytes and humoral and cellular immune responses, have been demonstrated. Also of great interest are the many studies suggesting their beneficial effects on age-associated changes in neurological health. [7] For instance, in vivo work revealed that PRPs alleviate beta-amyloid cytotoxicity in hippocampal neuronal cells; in humans, PRPs provided an early beneficial effect on cognitive symptoms and daily functioning at a dose as low as 100 mcg/d every other day. [8,9] Five grams of powder or eight capsules of IgG 2000 CWP provides 1% to 2% (50 to 100 mg) PRPs.\*

#### Oligosaccharides

Bovine milk oligosaccharides, which are a component of lactose, are modulators of gut microbiota. They provide protection by acting as decoys to attract antigens and inhibit them from binding to epithelial surfaces of the intestine. [10] Evidence also suggests that oligosaccharides act as growth promoters for a selected class of beneficial bacteria and contribute to the development and maturation of the intestinal immune response. [12] Approximately 66 acidic and neutral oligosaccharides, including sialyloligosaccharides, which are known for their high biological activity in humans, have been detected in IgG concentrate from colostral whey peptides.\*[13]

#### Gangliosides

Gangliosides are vital to the structure and function of cell membranes. They support neural development, and they have roles in supporting gut integrity, influencing immune cell signaling, modulating cytokine activity and production, and affecting the adherence and toxin production of antigens. [14-17] Research has demonstrated that providing gangliosides in the diet increases ganglioside content in the intestinal mucosa. Studies have indicated that low levels of gangliosides in the intestinal mucosa are associated with increased levels of cytokines, susceptibility to antigens, and poor gut integrity. [14] The ganglioside composition of bovine milk is predominately GM3 and GD3.\*[15]

Continued on next page

**Gastrointestinal Support** 

### IgG 2000 CWP™ Capsules Supplement Facts

Serving Size: 2 Capsules

	Amount Per Serving	%DV
Calories	5	
Protein	1 g	
lgG 2000 CWP™ (bovine-derived immunoglobulin concentrate) Immunoglobulin G (lgG)	1.25 g 500 mg	**
** Daily Value (DV) not established.		

Other Ingredients: HPMC (capsule), vegetable stearic acid, medium-chain triglycerides, vegetable magnesium stearate, and silica.
Contains: Milk.

**DIRECTIONS:** Take two capsules twice daily, or as directed by your healthcare practitioner.

Consult your healthcare practitioner prior to use. Individuals taking medication should discuss potential interactions with their healthcare practitioner. Do not use if tamper seal is damaged.

**DOES NOT CONTAIN:** Wheat, gluten, corn, yeast, soy, fish, shellfish, peanuts, tree nuts, egg, ingredients derived from genetically modified organisms (GMOs), artificial colors, artificial sweeteners, or preservatives.

**STORAGE:** Keep closed in a cool, dry place out of reach of children.



#### IgG 2000 CWP™ Powder Supplement Facts

Serving Size: 1 Tablespoon (about 5 g)

	Amount Per Serving	%DV
Calories	20	
Protein	4 g	
Calcium (from IgG 2000 CWP™)	40 mg	4%
IgG 2000 CWP™ (bovine-derived immunoglobulin concentrate) Immunoglobulin G (IgG)	5 g 2 g	**
** Daily Value (DV) not established.		

Other Ingredients: Sunflower lecithin. Contains: Milk.

**DIRECTIONS:** Mix one tablespoon (about 5 g) into 2-4 oz of water and consume daily, or take as directed by your healthcare practitioner.

Consult your healthcare practitioner prior to use. Individuals taking medication should discuss potential interactions with their healthcare practitioner. Do not use if tamper seal is damaged.

**DOES NOT CONTAIN:** Wheat, gluten, corn, yeast, soy, fish, shellfish, peanuts, tree nuts, egg, ingredients derived from genetically modified organisms (GMOs), artificial colors, artificial sweeteners, or preservatives.

**STORAGE:** Keep closed in a cool, dry place out of reach of children.

#### Chart: Bioactive Compounds in IgG 2000 CWP

Active	Key Function(s)
Immunoglobulins	Immune support*
Growth Factors	Lean mass support, cell and tissue repair and rejuvenation*
Sialic Acid	Immune modulation, brain health, prebiotic*
Lactoferrin	Immune support*
Proline-Rich Peptides (PRPs)	Immune modulation, brain and thymus support*
Oligosaccharides	Microbiota modulation*
Gangliosides	Immune cell signaling*

#### IgG 2000 CWP Studies

Building on the extensive literature pointing to the health benefits associated with bovine-derived colostrum and colostral whey, scientists performed in vitro, in vivo, and human clinical studies using IgG 2000 CWP to demonstrate its bioactivity and effectiveness. In vitro, IgG 2000 CWP was shown to support the body's normal repair pathways by stimulating the migration and production of gut epithelial cells.[18] In an in vivo model of gut integrity challenge, rats administered IgG 2000 CWP showed significant improvements in both microscopic and macroscopic mucosal health.[18] Supplemented dogs showed a significantly higher vaccine response and higher levels of fecal IgA when compared with the control group. These results were indicative of an enhanced immune status. Researchers also observed increases in gut microbiota diversity and stability in the supplemented dogs.[19] In humans (n = 12), 1000 mg/d of IgG 2000 CWP increased salivary IgA by 30%, improved quality of life scores, significantly increased the ability to perform activities of daily living, improved hyperimmune responses, and reduced minor health complaints by 47%. [20] In another human study (n = 7), a five-day challenge to gut integrity caused a three-fold rise from baseline in the ratio of lactulose to rhamnose (P < 0.01). This indicated increased gut permeability. Co-administration of IgG 2000 CWP inhibited this increase.\*[21]

#### References

- Godhia M, et al. Curr Res Nutr Food Sci. 2013;1(1):37-47. http://dx.doi. org/10.12944/CRNFSJ.1.1.04
- 2. Hurley D. [Dissertation]. Brookings, SD: South Dakota State University; 1994.
- **3.** Hurley WL, et al. *Nutrients*. 2011 Apr;3(4):442-74. [PMID: 22254105]
- **4.** Rump JA, et al. *Clin Investig*. 1992 Jul;70(7):588-94. [PMID: 1392428]
- **5.** Schaller JP, et al. *J Infect Dis.* 1992 Apr;165(4):623-30. [PMID: 1313067]
- 6. Berlutti F, et al. *Molecules*. 2011 Aug 16;16(8):6992-7018. [PMID: 21847071]
   7. Janusz M, et al. *Cell Mol Biol (Noisy-le-grand)*. 2013 Nov 3;59(1):4-11. [PMID:
- **8.** Froud KE, et al. *J Alzheimers Dis.* 2010;20(2):423-26. [PMID: 20164569]
- 9. Bilikiewicz A, et al. J *Alzheimers Dis.* 2004 Feb;6(1):17-26. [PMID: 15004324]
- Lane JA, et al. Int J Food Microbiol. 2012 Jul 2;157(2):182-8. [PMID: 22647676]
- **11.** Aldredge DL, et al. *Glycobiology*. 2013 Jun;23(6):664-76. [PMID: 23436288]
- 12. Lane JA, et al. Br J Nutr. 2013 Dec;110(12):2127-37. [PMID: 23710626]
- Barile D. Milk oligosaccharides. Confidential Report. Brookings, SD: Sterling Technology; 2011. [on file]
- **14.** Miklavcic JJ, et al. *J Nutr Metab*. 2012;2012:280286. [PMID: 22506104]
- **15.** Lee H, et al. *J Agric Food Chem.* 2013 Oct 9;61(40):9689-96. [PMID: 24024650]
- 16. Sánchez-Juanes F, et al. *Biol Chem.* 2009 Jan;390(1):31-40. [PMID: 18937626]
- Park EJ, et al. J Pediatr Gastroenterol Nutr. 2010 Mar;50(3):321-28. [PMID: 20118807]
- **18.** Playford R. [Unpublished colostral studies]. London, UK: Hammersmith Hospital;
- **19.** Satyaraj E, et al. *Br J Nutr.* 2013 Dec;110(12):2216-21. [PMID: 23773360]
- Jensen G, et al. Interim Report 9B. Brookings, SD: Sterling Technology. September 2010. [on file]
- 21. Playford RJ, et al. Clin Sci (Lond). 2001 Jun;100(6):627-33. [PMID: 11352778]

Additional references available upon request

#### EXCLUSIVE • PATENTEI