

RESVOXITROL



CLINICAL APPLICATIONS

- *Increases Antioxidant Protection*
- *Promotes a Balanced Inflammatory Response*
- *Activates Nrf2, the Master Cellular Switch for Antioxidant Production*
- *Supports Cellular Longevity*
- *Enhances Mechanisms of Cellular Detoxification*

CARDIOVASCULAR HEALTH

Resvoxitol is a unique combination of phytonutrients designed to enhance antioxidant potential by stimulating nuclear factor-erythroid-2-related factor 2 (Nrf2), a dynamic pathway known to increase the production of the body's most important cellular antioxidants. This unique combination of truebroc™, turmeric, pterostilbene, and resveratrol provides potent free radical scavenging potential, balancing of key inflammatory associated pathways, and induction of intracellular antioxidant production. These powerful phytochemicals are the basis of many "superfoods" and have extensive, peer-reviewed research supporting their dramatic effect on cell health and longevity.

Overview

Nrf2 is a transcription factor that regulates the antioxidant response of the body.¹ It is the primary cellular defense against the cytotoxic effects of oxidative stress.² This key protein is found inside a cell and is inactive until it is released by an Nrf2 activator. Once activated, Nrf2 migrates into the nucleus and binds to DNA at the location of the master antioxidant regulator, the antioxidant response element (ARE).³ Studies suggest Nrf2 plays an important role in supporting the activation of cellular antioxidant systems, as well as inflammatory balance.^{4,5} A number of fruit and vegetable compounds have been shown to activate Nrf2 and ARE, and increase antioxidant potential.⁶⁻⁹ It is believed that these compounds have a protective effect on the body, help maintain a healthy cycle of inflammation, and induce cellular longevity.¹⁰

Broccoli Seed Extract†

Broccoli seed extract has the highest levels of glucoraphanin (a potent Nrf2 stimulator) of any broccoli product available.¹¹ In fact, 230 mg of broccoli seed extract is equivalent to consuming

1 ¼ cups of fresh broccoli. Glucoraphanin is a phytochemical compound that acts as the precursor to sulforaphane, one of the leading compounds in broccoli that is responsible for many of its positive benefits.^{12,13} These compounds and their mechanisms have been studied extensively and supported by more than 500 scientific publications. Glucoraphanin and sulforaphane act as direct activators of Nrf2 and potent antioxidants shown to support detoxification and inflammatory balance.¹⁴⁻¹⁸ Studies show they work by improving hormone balance and increasing functional enzymes that support excretion of toxins from the body.¹⁹ Broccoli seed extract has been shown to remain active within the body for up to three days following consumption.²⁰

Turmeric Root Extract†

Turmeric is a bright orange-colored spice used for thousands of years in cooking and medicine. Through its antioxidant mechanisms, the active ingredient in turmeric, curcumin, helps to maintain a healthy cardiovascular system, support gastrointestinal (GI) health, and provide neuroprotective activity.²¹⁻²⁴ Curcumin has been shown to play a crucial role in maintaining proper inflammatory balance in a variety of bodily systems due to its ability to modulate Nrf2.²⁵ Studies have shown curcumin supports the maintenance of balanced inflammation in endothelial cells and cardiac cells.²⁶⁻²⁸ Curcumin has also been found to provide antioxidant protection to the cardiovascular system by reducing oxidative stress and supporting a normal immune response.²⁹

Pterostilbene†

Stilbenes make up a subgroup of phytochemicals that have potent antioxidant properties. Pterostilbene is a specific stilbene found in plant varieties like deerberry, rabbiteye

†These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

eFFICACY
the power of e

blueberries, unripe pinot noir grapes, *Botrytis vinifera*-infected Chardonnay grapes and Xarello grapes. Through activation of Nrf2, in vitro studies show pterostilbene significantly enhances expression of antioxidant enzymes, such as heme oxygenase-1 (HO-1) and glutathione reductase (GR).³⁰⁻³³ Some studies have shown antioxidant benefits may be improved when pterostilbene is combined with curcumin and resveratrol.^{34, 35}

Resveratrol†

Resveratrol, like pterostilbene, is a stilbene found in many plants and red wines. It is the most well-researched stilbene. Studies of resveratrol have found it to enhance the potential of antioxidant and detoxification activity through Nrf2 activation.³⁶ Other studies have linked resveratrol to enhancing blood sugar balance, balancing inflammation and reducing oxidative stress.^{37, 38}

Directions

2 capsules per day or as recommended by your health care professional.

Does Not Contain

Gluten, yeast, artificial colors and flavors.

Cautions

Do not consume this product if you are pregnant or nursing. Consult your physician for further information.

Supplement Facts ^{v2}		
Serving Size 2 Capsules		
Servings Per Container 30		
2 capsules contain	Amount Per Serving	% Daily Value
Turmeric Root Extract (Standardized to contain 95% Curcuminoids)	400 mg	*
Broccoli Seed Extract (truebroc™) (Standardized to contain 13% Glucoraphanin)	230 mg	*
trans-Resveratrol (from <i>Polygonum cuspidatum</i> (Roots))	100 mg	*
Pterostilbene (pTeroPure™)	60 mg	*
* Daily Value not established		

ID# 586060 60 Capsules

References

1. Nguyen T, Nioi P and Pickett CB. The Nrf2-antioxidant response element signaling pathway and its activation by oxidative stress. *Journal of Biological Chemistry*. 2009;284:13291-13295.
2. Cho J-M, Manandhar S, Lee H-R, Park H-M and Kwak M-K. Role of the Nrf2-antioxidant system in cytotoxicity mediated by anticancer cisplatin: implication to cancer cell resistance. *Cancer letters*. 2008;260:96-108.
3. Kensler TW, Wakabayashi N and Biswal S. Cell survival responses to environmental stresses via the Keap1-Nrf2-ARE pathway. *Annu Rev Pharmacol Toxicol*. 2007;47:89-116.
4. Chen X-L and Kunsch C. Induction of cytoprotective genes through Nrf2/antioxidant response element pathway: a new therapeutic approach for the treatment of inflammatory diseases. *Current pharmaceutical design*. 2004;10:879-891.
5. Li W, Khor TO, Xu C, Shen G, Jeong W-S, Yu S and Kong A-N. Activation of Nrf2-antioxidant signaling attenuates NFκB-inflammatory response and elicits apoptosis. *Biochemical pharmacology*. 2008;76:1485-1489.
6. Surh Y-J, Kundu JK and Na H-K. Nrf2 as a master redox switch in turning on the cellular signaling involved in the induction of cytoprotective genes by some chemopreventive phytochemicals. *Planta medica*. 2008;74:1526-1539.
7. Surh Y-J and Na H-K. NF-κB and Nrf2 as prime molecular targets for chemoprevention and cytoprotection with anti-inflammatory and antioxidant phytochemicals. *Genes & nutrition*. 2008;2:313-317.
8. Lee JH, Khor TO, Shu L, Su Z-Y, Fuentes F and Kong A-NT. Dietary phytochemicals and cancer prevention: Nrf2 signaling, epigenetics, and cell death mechanisms in blocking cancer initiation and progression. *Pharmacology & therapeutics*. 2013;137:153-171.
9. Gopalakrishnan A and Kong A-NT. Anticarcinogenesis by dietary phytochemicals: cytoprotection by Nrf2 in normal cells and cytotoxicity by modulation of transcription factors NF-κB and AP-1 in abnormal cancer cells. *Food and Chemical Toxicology*. 2008;46:1257-1270.

†These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

10. Su Z-Y, Shu L, Khor TO, Lee JH, Fuentes F and Kong A-NT. A perspective on dietary phytochemicals and cancer chemoprevention: oxidative stress, nrf2, and epigenomics. *Natural Products in Cancer Prevention and Therapy*: Springer; 2012: 133-162.
11. McWalter GK, Higgins LG, McLellan LI, Henderson CJ, Song L, Thornalley PJ, Itoh K, Yamamoto M and Hayes JD. Transcription factor Nrf2 is essential for induction of NAD (P) H: quinone oxidoreductase 1, glutathione S-transferases, and glutamate cysteine ligase by broccoli seeds and isothiocyanates. *The Journal of nutrition*. 2004;134:3499S-3506S.
12. Zhang Y, Kensler TW, Cho C-G, Posner GH and Talalay P. Anticarcinogenic activities of sulforaphane and structurally related synthetic norbornyl isothiocyanates. *Proceedings of the National Academy of Sciences*. 1994;91:3147-3150.
13. Clarke JD, Dashwood RH and Ho E. Multi-targeted prevention of cancer by sulforaphane. *Cancer letters*. 2008;269:291-304.
14. Singh AV, Xiao D, Lew KL, Dhir R and Singh SV. Sulforaphane induces caspase-mediated apoptosis in cultured PC-3 human prostate cancer cells and retards growth of PC-3 xenografts in vivo. *Carcinogenesis*. 2004;25:83-90.
15. Li Y, Zhang T, Korkaya H, Liu S, Lee H-F, Newman B, Yu Y, Clouthier SG, Schwartz SJ and Wicha MS. Sulforaphane, a dietary component of broccoli/broccoli sprouts, inhibits breast cancer stem cells. *Clinical Cancer Research*. 2010;16:2580-2590.
16. Heiss E, Herhaus C, Klimo K, Bartsch H and Gerhäuser C. Nuclear factor κ B is a molecular target for sulforaphane-mediated anti-inflammatory mechanisms. *Journal of Biological Chemistry*. 2001;276:32008-32015.
17. Thimmulappa RK, Mai KH, Srisuma S, Kensler TW, Yamamoto M and Biswal S. Identification of Nrf2-regulated genes induced by the chemopreventive agent sulforaphane by oligonucleotide microarray. *Cancer research*. 2002;62:5196-5203.
18. Morimitsu Y, Nakagawa Y, Hayashi K, Fujii H, Kumagai T, Nakamura Y, Osawa T, Horio F, Itoh K and Iida K. A sulforaphane analogue that potently activates the Nrf2-dependent detoxification pathway. *Journal of Biological Chemistry*. 2002;277:3456-3463.
19. Juge N, Mithen R and Traka M. Molecular basis for chemoprevention by sulforaphane: a comprehensive review. *Cellular and Molecular Life Sciences*. 2007;64:1105-1127.
20. Hanlon N, Coldham N, Gielbert A, Kuhnert N, Sauer MJ, King LJ and Ioannides C. Absolute bioavailability and dose-dependent pharmacokinetic behaviour of dietary doses of the chemopreventive isothiocyanate sulforaphane in rat. *British Journal of Nutrition*. 2008;99:559-564.
21. Nishinaka T, Ichijo Y, Ito M, Kimura M, Katsuyama M, Iwata K, Miura T, Terada T and Yabe-Nishimura C. Curcumin activates human glutathione S-transferase P1 expression through antioxidant response element. *Toxicology letters*. 2007;170:238-247.
22. Tapia E, Soto V, Ortiz-Vega KM, Zarco-Márquez G, Molina-Jijón E, Cristóbal-García M, Santamaría J, García-Niño WR, Correa F and Zazueta C. Curcumin induces Nrf2 nuclear translocation and prevents glomerular hypertension, hyperfiltration, oxidant stress, and the decrease in antioxidant enzymes in 5/6 nephrectomized rats. *Oxidative medicine and cellular longevity*. 2012;2012.
23. Aggarwal BB and Harikumar KB. Potential therapeutic effects of curcumin, the anti-inflammatory agent, against neurodegenerative, cardiovascular, pulmonary, metabolic, autoimmune and neoplastic diseases. *The international journal of biochemistry & cell biology*. 2009;41:40-59.
24. Zhang F, Altorki NK, Mestre JR, Subbaramaiah K and Dannenberg AJ. Curcumin inhibits cyclooxygenase-2 transcription in bile acid-and phorbol ester-treated human gastrointestinal epithelial cells. *Carcinogenesis*. 1999;20:445-451.
25. Balogun E, Hoque M, Pengfei G, Killeen E, Green CJ, Foresti R, Jawed A and Motterlini R. Curcumin activates the haem oxygenase-1 gene via regulation of Nrf2 and the antioxidant-responsive element. *Biochemical Journal*. 2003;371:887-895.
26. Motterlini R, Foresti R, Bassi R and Green CJ. Curcumin, an antioxidant and anti-inflammatory agent, induces heme oxygenase-1 and protects endothelial cells against oxidative stress. *Free Radical Biology and Medicine*. 2000;28:1303-1312.

27. Fiorillo C, Becatti M, Pensalfini A, Cecchi C, Lanzilao L, Donzelli G, Nassi N, Giannini L, Borchi E and Nassi P. Curcumin protects cardiac cells against ischemia-reperfusion injury: effects on oxidative stress, NF- κ B, and JNK pathways. *Free Radical Biology and Medicine*. 2008;45:839-846.
28. Yeh C-H, Chen T-P, Wu Y-C, Lin Y-M and Lin P-J. Inhibition of NF κ B Activation with Curcumin Attenuates Plasma Inflammatory Cytokines Surge and Cardiomyocytic Apoptosis Following Cardiac Ischemia/Reperfusion 1. *Journal of Surgical Research*. 2005;125:109-116.
29. Diaz MN, Frei B, Vita JA and Keaney Jr JF. Antioxidants and atherosclerotic heart disease. *New England Journal of Medicine*. 1997;337:408-416.
30. Chiou Y-S, Tsai M-L, Nagabhusanam K, Wang Y-J, Wu C-H, Ho C-T and Pan M-H. Pterostilbene is more potent than resveratrol in preventing azoxymethane (AOM)-induced colon tumorigenesis via activation of the NF-E2-related factor 2 (Nrf2)-mediated antioxidant signaling pathway. *Journal of agricultural and food chemistry*. 2011;59:2725-2733.
31. Saw CLL, Guo Y, Yang AY, Paredes-Gonzalez X, Ramirez C, Pung D and Kong A-NT. The berry constituents quercetin, kaempferol, and pterostilbene synergistically attenuate reactive oxygen species: involvement of the Nrf2-ARE signaling pathway. *Food and Chemical Toxicology*. 2014;72:303-311.
32. Yang Y, Wang J, Li Y, Fan C, Jiang S, Zhao L, Di S, Xin Z, Wang B and Wu G. HO-1 signaling activation by pterostilbene treatment attenuates mitochondrial oxidative damage induced by cerebral ischemia reperfusion injury. *Molecular neurobiology*. 2016;53:2339-2353.
33. McCormack D and McFadden D. Pterostilbene and cancer: current review. *Journal of Surgical Research*. 2012;173:e53-e61.
34. Matés J, Segura J, Alonso F and Márquez J. Natural antioxidants: therapeutic prospects for cancer and neurological diseases. *Mini reviews in medicinal chemistry*. 2009;9:1202-1214.
35. Liu RH. Potential synergy of phytochemicals in cancer prevention: mechanism of action. *The Journal of nutrition*. 2004;134:3479S-3485S.
36. Kode A, Rajendrasozhan S, Caito S, Yang S-R, Megson IL and Rahman I. Resveratrol induces glutathione synthesis by activation of Nrf2 and protects against cigarette smoke-mediated oxidative stress in human lung epithelial cells. *American Journal of Physiology-Lung Cellular and Molecular Physiology*. 2008;294:L478-L488.
37. Su H-C, Hung L-M and Chen J-K. Resveratrol, a red wine antioxidant, possesses an insulin-like effect in streptozotocin-induced diabetic rats. *American Journal of Physiology-Endocrinology and Metabolism*. 2006;290:E1339-E1346.
38. Martí AR, Villegas I, La Casa C and de la Lastra CA. Resveratrol, a polyphenol found in grapes, suppresses oxidative damage and stimulates apoptosis during early colonic inflammation in rats. *Biochemical pharmacology*. 2004;67:1399-1410.